

Neutron-Capture Elements in Halo, Thick-Disk, and Thin-Disk Stars: Neodymium

Mashonkina L., Kamaeva L., Samotoev V., Sakhibullin N.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

We have derived the LTE neodymium abundances in 60 cool stars with metallicities $[Fe/H]$ from 0.25 to -1.71 by applying a synthetic-spectrum analysis to spectroscopic observations of NdII lines with a resolution of $\lambda/\Delta\lambda \approx 60000$ and signal-to-noise ratios of 100-200. We have improved the atomic parameters of NdII and blending lines by analyzing the corresponding line profiles in the solar spectrum. Neodymium is overabundant with respect to iron in halo stars, $[Nd/Fe] = 0.33 \pm 0.09$, with the $[Nd/Fe]$ ratio decreasing systematically with metallicity when $[Fe/H] > -1$. This reflects an onset of efficient iron production in type I supernovae during the formation of the thick disk. The $[Nd/Ba]$ and $[Nd/Eu]$ abundance ratios behave differently in halo, thick-disk, and thin-disk stars. The observed abundance ratios in halo stars, $[Nd/Ba] = 0.34 \pm 0.08$ and $[Nd/Eu] = -0.27 \pm 0.05$, agree within the errors with the ratios of the elemental yields for the r-process. These results support the conclusion of other authors based on analyses of other elements that the r-process played the dominant role in the synthesis of heavy elements during the formation of the halo. The $[Nd/Ba]$ and $[Nd/Eu]$ ratios for thick-disk stars are almost independent of metallicity ($[Nd/Ba] = 0.28(\pm 0.03) - 0.01(\pm 0.04) [Fe/H]$ and $[Nd/Eu] = -0.13(\pm 0.03) + 0.05(\pm 0.04) [Fe/H]$) but are smaller in absolute value than the corresponding ratios for halo stars, suggesting that the synthesis of s-process nuclei started during the formation of the thick disk. The s-process is estimated to have contributed $\approx 30\%$ of the neodymium produced during this stage of the evolution of the Galaxy. The $[Nd/Ba]$ ratio decreases abruptly by 0.17 dex in the transition from the thick to the thin disk. The systematic decrease of $[Nd/Ba]$ and increase of $[Nd/Eu]$ with increasing metallicity of thin-disk stars point toward a dominant role of the s-process in the synthesis of heavy elements during this epoch. © 2004 MAIK "Nauka/Interperiodica".

<http://dx.doi.org/10.1134/1.1687012>
